
# **BRANDON** UNIVERSITY S. J. McKee Archives



## North Lauder locale Radiocarbon Report 3

### http://archives.brandonu.ca/en/permalink/descriptions12329

Part Of:	RG 7 Beverley Nicholson fonds
Description Level:	Sub sub series
Series Number:	2.5.3
Accession Number:	1-2010
GMD:	multiple media
Date Range:	1997-2000
Physical Description:	1 page
Material Details:	Radiocarbon date reports have been scanned in multi-page PDF files.
History / Biographical:	

North Lauder Radiocarbon Date report by IsoTrace Laboratory for Atkinson site #TO-13365.

#### Radiocarbon dating

The technique of radiocarbon dating was developed by Willard Libby and his colleagues at the University of Chicago in 1949.

Radiocarbon dating is used to estimate the age of organic remains from archaeological sites. Organic matter has a radioactive form of carbon (C14) that begins to decay upon death. C14 decays at a steady, known rate of a half life of 5,730 years. The technique is useful for material up to 50,000 years. Fluctuations of C14 in the atmosphere can affect results so dates are calibrated against dendrochronology. Radiocarbon dates are calibrated to calendar years.

Dates are reported in radiocarbon years or Before Present. Before Present refers to dates before 1950. The introduction of massive amounts of C14, due to atomic bomb and surface testing of atomic weapons, has widely increased the standard deviation on all dates after A.D. 1700 causing these dates to be unreliable.

Accelerated mass spectrometry can more accurately measure C14 with smaller samples and can date materials to 80,000 years.

Scope and Content:

Sub sub series contains radiocarbon dates from: Atkinson site and Flintstone Hill.

Name Access:	North Lauder locale Radiocarbon Report 3
Subject Access:	Archaeology
	North Lauder locale
	North Lauder locale Radiocarbon Report 3

### Documents

Athins	100 LINUTANA ANA COLUMN TI AND
	Output by calibration program dischole Oppyright (c) R.B.Beubens
	12-24-03
	1338 In 1-8-3 binne - siner enqui
	incarios daim - 1000 j 120 MP arbitisms, with a perdudtific of 154 or presider for the calibrated age
15 645 544 140	this reducentee data, have been calculated from the dealers salibation as the 6H and 9H conditioner intervals, which are the in and it intro- a scenari distribution, are the given, a probability of 1000 means the location data intervalsment the dealer calculation reveares with
	reading and monohol to the assument multiple of 5 years. . Probability set Age 00.3 % c.1. 00.5 % c.1.
	the b mant call and plant and a state and and a state and
1	
121 19 7,	illusted with the standard data on $form$ , there is a standard data of the standard data and the standard data of the standard data o
100 W	0.4
80 70	
8	
stability 5 \$ \$	P
- M-	
	N VL
1	000 0708 000 000 000 000 000 000 000
(11	K) Galilentel ap
0 E -	Attingen DC14
Z.O.,	3 Atkinson RC14
	TO-13365.pdf
	10-15505.pul
	Read PDF
1	Download PDF
4	Download PDF