# **BRANDON** UNIVERSITY S. J. McKee Archives

beTrace Radiocarbon Laboratory	the billing face. Treasure states of the Canada Mill 1477
Acutorator Marc Spectrometry Robby at the Entransity of Thomas	Matter (0:10) 401 Fe(-0)-95-85 Feat self-sense-boost-co
Radman ban sandari	
stag of, sea	
laborate di Materiano dependitativo desilore di antere di	dr. Results 148
The second is the section of the spaces whethere benefit points single the statement is solved as yell in the transmission of the section and should a simulation of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the the section of the section of the section of the the section of the section of the section of the section of the section of the section of the section of the the section of the secti	annel "Coffe and the output age is good to provide the state the output of the second second se
l nast ik o kor ya nastada nasta kuali if No ini Di ya nast ani an nastajik	Objection of the second s

## North Lauder locale Radiocarbon Report 2

### http://archives.brandonu.ca/en/permalink/descriptions12328

Part Of:	RG 7 Beverley Nicholson fonds
Description Level:	Sub sub series
Series Number:	2.5.2
Accession Number:	1-2010
GMD:	multiple media
Date Range:	1997-2000
Physical Description:	2 pages
Material Details:	Radiocarbon date reports have been scanned in multi-page PDF files.
History / Biographical:	
Biographical:	

North Lauder Radiocarbon Date report by IsoTrace Laboratory for Atkinson site #TO-10640.

#### Radiocarbon dating

The technique of radiocarbon dating was developed by Willard Libby and his colleagues at the University of Chicago in 1949.

Radiocarbon dating is used to estimate the age of organic remains from archaeological sites. Organic matter has a radioactive form of carbon (C14) that begins to decay upon death. C14 decays at a steady, known rate of a half life of 5,730 years. The technique is useful for material up to 50,000 years. Fluctuations of C14 in the atmosphere can affect results so dates are calibrated against dendrochronology. Radiocarbon dates are calibrated to calendar years.

Dates are reported in radiocarbon years or Before Present. Before Present refers to dates before 1950. The introduction of massive amounts of C14, due to atomic bomb and surface testing of atomic weapons, has widely increased the standard deviation on all dates after A.D. 1700 causing these dates to be unreliable.

Accelerated mass spectrometry can more accurately measure C14 with smaller samples and can date materials to 80,000 years.

Scope and Content:

Sub sub series contains radiocarbon dates from: Atkinson site and Flintstone Hill.

Name Access:	North Lauder locale Radiocarbon Report 2
Subject Access:	Archaeology
	North Lauder locale
	North Lauder locale Radiocarbon Report 2

#### Documents

	IsoTrace Radiocarbon Laboratory	60 St. George Street Terrenie (Orie) George Mill 147
	Accelerator Mass Spectroserby Facility active University of Torenta	Telephone 414 - 575 - 4620 Pere 418 - 576 - 4751 Tanali mell'Indone Parlamente
	Badiocarbas Analysis Ba	uport.
	May 27, 2008	
	Sobscribe: 30 Nuberlaws, Dept of Native Studies, Rouslin Unit, F. This small is the average of 2 separate analysis (control provider) integra functionation to a loss of $D^{-1}C_{2} = 15\%$ , using the reasons on an avoid-based conversion of advanced on this into your before of 9075 years. The series represents the 68.3 % confidence lasts.	and is consisted for second and quittering of 752/7C rates. The second second control
	Sample Weight Matelification Discription used (ag TS-1 Oldetroor Reark): elutered http: 544	belliner Apr Labrandher (ywan BP) TG-10640 5250 ± 00
	) result like to how your constants on this result. If this result is to if you could send up a reprint.	and in a publication, I would approvate
		The.
		DETAM
		Dr. E. F. Besters
2	5.2 Atkinso	n RC14
	_	
	TO-10640	).pdf
		o.p.a.
	<b>P</b> Road	DDE
	🗐 Read	PDF
	Read	